

PATIENT NAME: DOE, JOHN	DATE OF BIRTH: June 02, 1942	GENDER: MALE	HEIGHT: 5 ft 9 inch	WEIGHT: 140 lb	BMI: 20.5
-----------------------------------	--	------------------------	-------------------------------	--------------------------	---------------------


NAME OF PHYSICIAN: DR. JAMES DOE	DATE OF TEST TAKEN: October 25, 2020
---	---

BIOMARKER RESULTS

Sample Date	October 19, 2020		
alpha-2-Macroglobulin	366 mg/dL	ALT	15 IU/L
Haptoglobin	68 mg/dL	AST	17 IU/L
Apolipoprotein A1	152 mg/dL	Fasting Glucose	87 mg/dL
Total Bilirubin	0.585 mg/dL	Total Cholesterol	157 mg/dL
GGT	22 IU/L	Triglycerides	98 mg/dL

SCORES

FIBROSIS




F3
Significant Fibrosis
0.73

FIBROSIS:

Liver fibrosis represents scarring of the liver. Scarring of the liver is due to reaction from viral infections, fat and/or alcohol.

ACTIVITY




A0
No Activity
0.09

ACTIVITY:

Liver activity represents inflammation of the liver. Inflammation of the liver is due to reaction from viral infections, fat and/or alcohol.

STEATOSIS



S0
No Steatosis
0.11

STEATOSIS:

Liver steatosis is the accumulation of fat in the liver. Liver steatosis is commonly due to metabolic anomalies (e.g. Dyslipidemia, diabetes, overweight/obesity) and/or alcohol.

INTERPRETATION

- Your result for the SAF score is S0-A0-F3.
- This score indicates that you have no steatosis, no inflammatory activity and significant fibrosis.
- Consult your physician for further evaluation.

PRECAUTIONS:

- The reliability of results is dependent on compliance with the preanalytical and analytical conditions recommended by Fibronostics.
- The test must be deferred for acute hemolysis, acute hepatitis, acute inflammation, extra hepatic cholestasis.
- The advice of a specialist must be sought for interpretation in chronic hemolysis and Gilbert's syndrome.
- The test interpretation is not validated for liver transplant patients.
- Isolated extreme values of any of the biomarker results should lead to caution in interpreting the results.
- In the case of discordance between a biopsy result and a test, it is recommended to seek the advice of a specialist.
- The causes of these discordances could be due to a flaw of the test or to a flaw in the biopsy: i.e. a liver biopsy has a 33% variability rate for one fibrosis stage.
- The fibrosis score is interpretable for chronic hepatitis B, chronic hepatitis C, alcoholic steatosis and non-alcoholic steatosis.
- The activity score is interpretable for chronic hepatitis B and chronic hepatitis C.

Director:
Imtiaz Alam, MD

This report is computer generated. No signature required.

CLIA # 10D2190357